

# Effective Prime Uniqueness

Peter Cholak and Charlie McCoy

VITA CEDO  
June, 2015

Germany

# Prime Models

## Definition

A countable model  $\mathcal{A}$  is *prime* if for all models  $\mathcal{B}$  if  $\mathcal{A} \equiv \mathcal{B}$  then  $\mathcal{A} \preceq \mathcal{B}$ .

- Let  $T$  be the first order theory of  $\mathcal{A}$ . So  $\mathcal{A}$  elementary embeds into every model  $\mathcal{B}$  of  $T$ .
- $T$  will always be a complete first order theory in a countable language.
- We want to explore the notion of prime structures. First classically then effectively.

# Types

## Definition

Given a model  $\mathcal{A}$  and a tuple  $\vec{a} \in |\mathcal{A}|$  the *type* of  $a$ ,  $p(\vec{a})$  is the theory of  $(\mathcal{A}, \vec{a})$ .

## Lemma

*If  $\mathcal{A} \preccurlyeq \mathcal{B}$  via  $f$  then, for all  $\vec{a} \in |\mathcal{A}|$ , the types of  $\vec{a}$  and  $f(\vec{a})$  are the same.*

## Definition

A type  $p(\vec{a})$  is principal iff there is a formula  $\phi(\vec{a})$  such that, for all  $\sigma(\vec{a}) \in p(\vec{a})$ ,  $T \vdash \phi(\vec{a}) \rightarrow \sigma(\vec{a})$ . ( $\phi$  is called an atom of  $T$  and the complete formula of  $p(\vec{a})$ .)

# Atomic Models

## Theorem (Omitting Types)

*If a type  $p(\vec{a})$  is nonprincipal there is a model  $\mathcal{A}$  omitting it.*

## Definition

A model  $\mathcal{A}$  is atomic if all its types are principal.

## Corollary

*All prime models are atomic.*

# First Isomorphism Result

## Lemma

*If  $\mathcal{A}$  and  $\mathcal{B}$  are atomic models of  $T$  then they are isomorphic.*

## Proof.

Build the isomorphism  $f$  stagewise using a back and forth argument making sure the types of the domain and range remain the same.

(Forth) Assume that  $f_{2s}$  is a partial finite function with domain  $\vec{a}$  such that the types of  $\vec{a}$  and  $f_{2s}(\vec{a})$  are the same. At stage  $2s + 1$ , let  $d$  be the first element of  $\mathcal{A}$  not in  $\vec{a}$ , **let  $\phi(\vec{a}, d)$  be the complete formula of  $p(\vec{a}, d)$** , then let  $f_{2s+1}(d)$  be the first element of  $\mathcal{B}$  not in  $f_{2s}(\vec{a})$  such that  $\mathcal{B} \models \phi(f_{2s}(\vec{a}), f_{2s+1}(d))$ .

(Back) Similar.



# Two Useful Corollaries from the Isomorphism Result

## Corollary

*Atomic models are prime.*

## Corollary (Prime Uniqueness)

*Two prime models are isomorphic.*

# Complete formulas

## Lemma

$\{\phi \mid \phi \text{ is a complete formula}\}$  is  $\Pi_1^T$ .

## Proof.

Check all  $\sigma$  does  $T \vdash \phi \rightarrow \sigma$  or  $T \vdash \phi \rightarrow \neg\sigma$ . □

## Lemma (Folklore)

*There are computable complete theories  $T$  ( $T$  is called decidable) where  $\{\phi \mid \phi \text{ is a complete formula}\}$  is  $\Pi_1^0$ -complete.*

## Sketch.

Use unary predicates  $R_i$  and  $R_{i,s}$  to model  $\phi_i(i)$  and  $\phi_{i,s}(i)$ . □

There are also decidable  $T$  where the complete formulas are computable. This is also dependent on the language.

# Atomic Theories

## Definition

A theory  $T$  is *atomic* if for every formula  $\sigma$  there is an atom  $\phi$  of  $T$  such that  $T \vdash \phi \rightarrow \sigma$ .

## Lemma (AMT)

*Every atomic theory has an atomic model.*

AMT was explored by Hirschfeldt, Shore and Slaman. It turns out that this does not hold effectively, is incomparable to WKL and is properly below ACA.

## Question

*Does prime uniqueness hold effectively?*

Yes. Hence this talk and corresponding paper.



# Effectively Prime and Atomic Models

## Definition

Let  $T$  be a decidable theory and  $\mathcal{A}$  a decidable model of  $T$ .

- The model  $\mathcal{A}$  is *effectively prime*, if for every decidable model  $\mathcal{M} \models T$ , there is a computable elementary embedding  $f : \mathcal{A} \rightarrow \mathcal{M}$ . Note that  $f$  need not be uniformly computable in  $\mathcal{A}$  and/or  $\mathcal{M}$ .
- The model  $\mathcal{A}$  is *effectively atomic* if there is a computable function  $g$  that accepts as an input a tuple  $\vec{a}$  from  $\mathcal{A}$  (of any length) and outputs a complete formula  $\varphi(\vec{x})$  so that  $\mathcal{A} \models \varphi(\vec{a})$ . Again  $g$  need not be uniformly computable in  $\mathcal{A}$ .
- The model  $\mathcal{A}$  is *uniformly effectively prime* if there is a partial computable function  $\Phi$  so that, given  $\mathcal{M} \models T$ ,  $\Phi(\mathcal{M})$  halts and outputs the code of a computable elementary embedding  $f : \mathcal{A} \rightarrow \mathcal{M}$ . Again  $\Phi$  need not be uniformly computable in  $\mathcal{A}$ .

# Effective Corollaries from the Isomorphism Result

## Corollary

*If two decidable models  $\mathcal{A}$  and  $\mathcal{B}$  of the same decidable theory  $T$  are both effectively atomic, then the classical back and forth construction produces a computable isomorphism between  $\mathcal{A}$  and  $\mathcal{B}$ .*

## Corollary

*Effectively atomic implies uniformly effectively prime.*

Effectively prime = effectively atomic = uniformly  
effectively prime

### Theorem ( $\text{RCA}_0$ )

*There is a Turing functional  $\Phi(\mathcal{A}, e)$  such that if  $T$  is decidable and  $\mathcal{A} \models T$  is a decidable model then either, for some  $e$ ,  $\Phi(\mathcal{A}, e)$  witnesses that  $\mathcal{A}$  is effectively atomic or there is a decidable  $\mathcal{M} \models T$ , such that there is no computable elementary embedding of  $\mathcal{A}$  into  $\mathcal{M}$ .*

## Cannot be improved!

### Lemma (Folklore)

*For all  $\Phi$ , there is an effectively atomic  $\mathcal{A}$  such that  $\Phi(\mathcal{A})$  does not witness that  $\mathcal{A}$  is effectively atomic.*

### Sketch.

Use the recursion theorem for a code of  $\mathcal{A}$ . Work in the language of infinitely many unary relations and depending on  $\Phi(\mathcal{A})$  the resulting model has nothing in any of these relations or exactly one of the relations splits the model into 2 infinite parts. □

Hence the “obvious” notion of “uniformly effectively atomic” is vacuous. Again this is also dependent on the language and  $T$ . But note a code for  $\mathcal{A}$  computes the theory  $T$ .

## A Preliminary Result

### Theorem

*Let  $T$  be decidable and  $\mathcal{A}, \mathcal{B} \models T$  be decidable models. Then either there is a computable isomorphism  $h : \mathcal{A} \cong \mathcal{B}$ ; or there is a decidable  $\mathcal{M} \models T$ , so that either there is no computable elementary embedding of  $\mathcal{A}$  into  $\mathcal{M}$ , or there is no computable elementary embedding of  $\mathcal{B}$  into  $\mathcal{M}$ .*

## The Construction

Given  $\mathcal{A}$ . Build  $\mathcal{M}$  via a Henkin construction using a finite priority argument to meet the following:

$\mathcal{R}_\Psi$  :  $\neg(\mathcal{A} \preceq \mathcal{M} \text{ via } \Psi)$  or there is a  $g$  witnessing  
that  $\mathcal{A}$  is effectively atomic.

If  $\Psi$  is a permutation then we meet  $\mathcal{R}_\Psi$  via different types, for some  $\vec{a}$ , the types of  $\vec{a}$  and  $\Psi(\vec{a})$  are different. Meeting  $\mathcal{R}_\Psi$  is  $\Sigma_2$ . When adding formulas  $\sigma_s$  for the diagram of  $\mathcal{M}$  one *carefully* looks for ways to diagonalize for some  $\vec{a}$  and  $\Psi(\vec{a})$ . The failure to diagonalize for some permutation  $\Psi$  produces  $g$ .

## In $RCA_0$

### Proof.

First at each stage use  $\Sigma_1$  induction to show that there is no requirement or a least requirement for which we can diagonalize.

Meeting  $\mathcal{R}_\Psi$  is  $\Sigma_2$ : either is not total ( $\Sigma_2$ ), is not onto ( $\Sigma_2$ ), is not 1 – 1 ( $\Sigma_1$ ), or, for some  $\vec{a}$ , the types of  $\vec{a}$  and  $\Psi(\vec{a})$  are different ( $\Sigma_1$ ). Assume that  $\Psi$  is an embedding of  $\mathcal{A}$  into  $\mathcal{M}$  then there by  $\Sigma_1$  induction is a stage where all higher priority requirements stop acting (impact how  $\sigma_s$  is added to the diagram of  $\mathcal{M}$ ). So after that stage if we can diagonalize to beat  $\Psi$  we will. This allows us to show  $\mathcal{A}$  is effectively atomic. □